

Desulfomicrobium baculatum [NiFeSe]-hydrogenase



References:

1. E Garcin, X Vernede, EC Hatchikian, A Volbeda, M Frey & JC Fontecilla-Camps. The crystal structure of a reduced [NiFeSe] hydrogenase provides an image of the activated catalytic center. *Structure* (1999) **7**, 557-566.
2. A Parkin, G Goldet, C Cavazza, JC Fontecilla-Camps & FA Armstrong. The difference a Se makes? Oxygen-tolerant hydrogen production by the [NiFeSe]-hydrogenase from *Desulfomicrobium baculatum*. *J Am Chem Soc* (2008) **130**, 13410-13416.
3. E Reisner, JC Fontecilla-Camps & FA Armstrong. Catalytic electrochemistry of a [NiFeSe]-hydrogenase on TiO₂ and demonstration of its suitability for visible-light driven H₂ production. *Chem Commun* (2009) **5**, 550-552.
4. E Reisner, DJ Powell, C Cavazza, JC Fontecilla-Camps & FA Armstrong. Visible light-driven H₂ production by hydrogenases attached to dye-sensitized TiO₂ nanoparticles. *J Am Chem Soc* (2009) **131**, 18457-18466.
5. A Volbeda, P Amara, M Iannello, AL DeLacey, C Cavazza & JC Fontecilla-Camps. Structural foundations for the O₂-resistance of *Desulfomicrobium baculatum* [NiFeSe]-hydrogenase, *Chem Comm* (2013) **49**, 7061-7063