




Desulfovibrio gigas [NiFe]-hydrogenase




Location of metal sites at 5 Å resolution¹



First crystal structure refined at 2.8 Å resolution²



Evidence for CO and CN ligands bound to Fe^{3,4}



Detection of tunnels at 2.54 Å resolution⁵

References:

1. A Volbeda, C Piras, MH Charon, EC Hatchikian, M Frey & JC Fontecilla-Camps. Location of the redox centers in hydrogenase as determined by X-ray crystallography at 5 Å resolution. *ESF/CCP4 Newsletter* (1993) **28**, 30-33.
2. A Volbeda, MH Charon, C Piras, EC Hatchikian, M Frey & JC Fontecilla-Camps. Crystal Structure of the Nickel-Iron Hydrogenase from *Desulfovibrio gigas*. *Nature* (1995) **373**, 580-587.
3. A Volbeda, E Garcin, C Piras, AL de Lacey, VM Fernandez, EC Hatchikian, M Frey & JC Fontecilla-Camps. Structure of the [NiFe] hydrogenase active site: evidence for biologically uncommon Fe ligands. *J Am Chem Soc* (1996) **118**, 12989-12996.
4. AL de Lacey, EC Hatchikian, A Volbeda, M Frey, JC Fontecilla-Camps & VM Fernandez. Infrared-spectroelectrochemical characterization of the [NiFe] hydrogenase of *Desulfovibrio gigas*. *J Am Chem Soc* (1997) **119**, 7181-7189.
5. Y Montet, P Amara, A Volbeda, X Vernede, EC Hatchikian, MJ Field, M Frey & JC Fontecilla-Camps. Gas access to the active site of Ni-Fe hydrogenases probed by X-ray crystallography and molecular dynamics. *Nature Struct Biol* (1997) **4**, 523-526.
6. P Amara, A Volbeda, JC Fontecilla-Camps & MJ Field. A hybrid density functional theory / molecular mechanics study of nickel-iron hydrogenase: investigation of the active site redox states. *J Am Chem Soc* (1999) **121**, 4468-4477.
7. A Volbeda & JC Fontecilla-Camps: The active site and catalytic mechanism of NiFe hydrogenases. *Dalton transactions* (2003), **21**, 4030-4038.